

ENVIRONMENTAL PROTECTION



NRA

*National Rivers Authority
South West Region*

**River Looe Catchment
River Water Quality
Classification 1990**

**NOVEMBER 1991
WQP/91/016
B L MILFORD**

**GORDON H BIELBY BSc
Regional General Manager**

**C V M Davies
Environmental Protection
Manager**

ACKNOWLEDGEMENTS

The Water Quality Planner acknowledges the substantial contributions made by the following staff:

- R. Broome - Co-ordinator and Editor
- A. Burrows - Production of Maps and editorial support
- P. Grigorey - Production of Maps and editorial support
- B. Steele - Production of Forepage
- C. McCarthy - Administration and report compilation

Special thanks are extended to A. Burghes of Moonsoft, Exeter for computer support and the production of statistical schedules.

The following NRA sections also made valuable contributions:

- Pollution Control
- Field Control and Wardens
- Water Resources

Thanks also to R. Hamilton and J. Murray-Bligh for their contributions.

Suggestions for improvements that could be incorporated in the production of the next Classification report would be welcomed.

Further enquiries regarding the content of these reports should be addressed to:

Freshwater Scientist,
National Rivers Authority,
Manley House,
Kestrel Way,
EXETER,
Devon EX2 7LQ

ENVIRONMENT AGENCY



129238

RIVER WATER QUALITY IN THE RIVER LOOE CATCHMENT

LIST OF CONTENTS

	Page No.
1 Introduction	1
2 River Looe Catchment	1
3 National Water Council's River Classification System	2
4 1990 River Water Quality Survey	3
5 1990 River Water Quality Classification	3
6 Non-compliance with Quality Objectives	4
7 Causes of Non-compliance	4
8 Glossary of Terms	5
9 References	5
10 Appendices:	
10.1 River Quality Objectives including Monitoring points	
10.2 Basic Determinand Analytical Suite	
10.3 National Water Council (NWC) River Classification System	
10.4 NWC Criteria for Non-Metallic Determinands - Regional Variation	
10.4.1 NWC Criteria for Metallic Determinands - Regional Variation	
10.5 1990 River Water Quality Classification - tabular format	
10.6 1990 River Water Quality Classification - map format	
10.7 Calculated Determinand Statistics used for Quality Assessment	
10.8 Compliant/Non-Compliant River Reaches	
10.9 Number of Samples Results exceeding quality standards	
10.10 Percentage Exceedance of Determinand Statistics from Quality Standard	
10.11 Identification of Possible Causes of Non-Compliance with River Quality Objectives	

National Rivers Authority South West Region



NRA

National Rivers Authority
South West Region


NRA Wessex Region

EXETER

BODMIN

**River Looe
Catchment**

River Looe Catchment



1. INTRODUCTION

Monitoring to assess the quality of river waters is undertaken in thirty-two catchments within the region. As part of this monitoring programme samples are collected routinely from selected monitoring points at a pre-determined frequency per year, usually twelve spaced at monthly intervals. Each monitoring point provides data for the water quality of a river reach (in kilometres) upstream of the monitoring point.

River lengths have been re-measured and variations exist over those recorded previously.

Each water sample collected from each monitoring point is analysed for a range of chemical and physical constituents or properties known as determinands. The analytical results for each sample are entered into a computer database called the Water Quality Archive.

Selected data are accessed from the Archive so that the quality of each river reach can be determined based on a River Classification System developed by the National Water Council (NWC), (9.1).

This report presents the river water quality classification for 1990 for monitored river reaches in the River Looe catchment.

2. RIVER LOOE CATCHMENT

The East Looe River flows over a distance of 12.8 km from its source to the tidal limit, (Appendix 10.1). Water quality was monitored at seven locations on the main river; six of these sites were sampled at approximately monthly intervals. The site downstream of Liskeard sewage treatment works was sampled on twenty occasions during 1990 because of no recent water quality data.

The West Looe River flows over a distance of 12.1 km from its source to the tidal limit, (Appendix 10.1). Water Quality was monitored at four locations on the main river; three of these sites were sampled at approximately monthly intervals and the site at Bosent Bridge was sampled on twenty occasions during 1990 because of no recent water quality data.

Throughout the East Looe catchment one secondary tributary of the East Looe River was sampled at approximately monthly intervals.

Throughout the West Looe catchment one secondary tributary of the West Looe River was sampled on twenty occasions during 1990 because of no recent water quality data and one other secondary tributary was sampled at approximately monthly intervals.

Polperro Stream flows over a distance of 7.0 km from its source to the tidal limit, (Appendix 10.1) and was monitored at one location situated in the lower reaches of the stream.

2.1 SECONDARY TRIBUTARIES

The Connon Stream flows over a distance of 5.3 km from its source to the confluence with the West Looe River, (Appendix 10.1) and was monitored at three locations.

The Coldrinnick Stream flows over a distance of 5 km from its source to the confluence with the West Looe River, (Appendix 10.1) and was monitored at one location situated in the lower reaches of the stream.

The Dobwalls Stream flows over a distance of 2.2 km from its source to the confluence with the East Looe River, (Appendix 10.1) and was monitored at one location situated in the lower reaches of the stream.

Each sample was analysed for a minimum number of determinands (Appendix 10.2) plus additional determinands based on local knowledge of the catchment. In addition, at selected sites, certain metal analyses were carried out.

The analytical results from all of these samples have been entered into the Water Quality Archive and can be accessed through the Water Act Register, (9.2).

3. NATIONAL WATER COUNCIL'S RIVER CLASSIFICATION SYSTEM

3.1 River Quality Objectives

In 1978 river quality objectives (RQOs) were assigned to all river lengths that were part of the routine monitoring network and to those additional watercourses, which were not part of the routine network, but which received discharges of effluents.

For the majority of watercourses long term objectives were identified based on existing and assumed adequate quality for the long term protection of the watercourse. In a few instances short term objectives were identified but no timetable for the achievement of the associated long term objective was set.

The RQOs currently in use in the River Looe catchment are identified in Appendix 10.1.

3.2 River Quality Classification

River water quality is classified using the National Water Council's (NWC) River Classification System (see Appendix 10.3), which identifies river water quality as being one of five quality classes as shown in Table 1 below:

Table 1 - National Water Council - River Classification System

<u>Class</u>	<u>Description</u>
1A	Good quality
1B	Lesser good quality
2	Fair quality
3	Poor quality
4	Bad quality

Using the NWC system, the classification of river water quality is based on the values of certain determinands as arithmetic means or as 95 percentiles (5 percentiles are used for pH and dissolved oxygen) as indicated in Appendices 10.4.1 and 10.4.2.

The quality classification system incorporates some of the European Inland Fisheries Advisory Commission (EIFAC) criteria (Appendix 10.3) recommended for use by the NWC system.

4. 1990 RIVER WATER QUALITY SURVEY

The 1990 regional classification of river water quality also includes the requirements of the Department of the Environment quinquennial national river quality survey. The objectives for the Department of the Environment 1990 River Quality Survey are given below:

- 1) To carry out a National Classification Survey based on procedures used in the 1985 National Classification Survey, including all regional differences.
- 2) To classify all rivers and canals included in the 1985 National Classification Survey.
- 3) To compare the 1990 Classification with those obtained in 1985.

In addition, those watercourses, which were not part of the 1985 Survey and have been monitored since that date, are included in the 1990 regional classification of river water quality.

5. 1990 RIVER WATER QUALITY CLASSIFICATION

Analytical data collected from monitoring during 1988, 1989 and 1990 were processed through a computerised river water quality classification programme. This resulted in a quality class being assigned to each monitored river reach as indicated in Appendix 10.5.

The quality class for 1990 can be compared against the appropriate River Quality Objective and previous annual quality classes (1985-1989) also based on three years combined data, for each river reach in Appendix 10.5.

The river water classification system used to classify each river length is identical to the system used in 1985 for the Department of the Environment's 1985 River Quality Survey. The determinand classification criteria used to determine the annual quality classes in 1985, subsequent years and for 1990 are indicated in Appendices 10.4 and 10.4.1.

Improvements to this classification system could have been made, particularly in the use of a different suspended solids standard for Class 2 waters. As the National Rivers Authority will be proposing new classification systems to the Secretary of State in the near future, it was decided to classify river lengths in 1990 with the classification used for the 1985-1989 classification period.

The adoption of the revised criteria for suspended solids in Class 2 waters would not have affected the classification of river reaches.

The river quality classes for 1990 of monitored river reaches in the catchment are shown in map form in Appendix 10.6.

The calculated determinand statistics for pH, temperature, dissolved oxygen, biochemical oxygen demand (BOD), total ammonia, un-ionised ammonia, suspended solids, copper and zinc from which the quality class was determined for each river reach, are indicated in Appendix 10.7.

6. NON-COMPLIANCE WITH QUALITY OBJECTIVES

Those monitored river reaches within the catchment, which do not comply with their assigned (RQO), are shown in map form in Appendix 10.8.

Appendix 10.9 indicates the number of samples analysed for each determinand over the period 1988 to 1990 and the number of sample results per determinand, which exceed the determinand quality standard.

For those non-compliant river reaches in the catchment, the extent of exceedance of the calculated determinand statistic with relevant quality standard (represented as a percentage), is indicated in Appendix 10.10.

7. CAUSES OF NON-COMPLIANCE

For those river reaches, which did not comply with their assigned RQOs, the cause of non-compliance (where possible to identify) is indicated in Appendix 10.11.

8. GLOSSARY OF TERMS


RIVER REACH	A segment of water, upstream from sampling point to the next sampling point.
RIVER LENGTH	River distance in kilometres.
RIVER QUALITY OBJECTIVE	That NWC class, which protects the most sensitive use of the water.
95 percentiles	Maximum limits, which must be met for at least 95% of the time.
5 percentiles	Minimum limits, which must be met for at least 95% of the time.
BIOLOGICAL OXYGEN DEMAND (5 day carbonaceous ATU)	A standard test measuring the microbial uptake of oxygen - an estimate of organic pollution.
pH	A scale of acid to alkali.
UN-IONISED AMMONIA	Fraction of ammonia poisonous to fish, NH^3 .
SUSPENDED SOLIDS	Solids removed by filtration or centrifuge under specific conditions.
USER REFERENCE NUMBER	Reference number allocated to a sampling point.
INFERRED STRETCH	Segment of water, which is not monitored and whose water quality classification is assigned from the monitored reach upstream.

9. REFERENCES

Reference

- 9.1 National Water Council (1977). River Water Quality: The Next Stage. Review of Discharge Consent Conditions. London.
- 9.2 Water Act 1989 Section 117
- 9.3 Alabaster J. S. and Lloyd R. Water Quality Criteria for Freshwater Fish, 2nd edition, 1982. Butterworths.

Looe Catchment River Quality Objectives



BASIC DETERMINAND ANALYTICAL SUITE FOR ALL CLASSIFIED RIVER SITES

pH as pH Units

Conductivity at 20 C as uS/cm

Water temperature (Cel)

Oxygen dissolved & saturation

Oxygen dissolved as mg/l O

Biochemical oxygen demand (5 day total ATU) as mg/l O

Total organic carbon as mg/l C

Nitrogen ammoniacal as mg/l N

Ammonia un-ionised as mg/l N

Nitrate as mg/l N

Nitrite as mg/l N

Suspended solids at 105 C as mg/l

Total hardness as mg/l CaCO₃

Chloride as mg/l Cl

Orthophosphate (total) as mg/l P

Silicate reactive dissolved as mg/l SiO₂

Sulphate (dissolved) as mg/l SO₄

Sodium (total) as mg/l Na

Potassium (total) as mg/l K

Magnesium (total) as mg/l Mg

Calcium (total) as mg/l Ca

Alkalinity as pH 4.5 as mg/l CaCO₃

NWC RIVER QUALITY CLASSIFICATION SYSTEM

River Class	Quality criteria	Remarks	Current potential uses
	Class limiting criteria (95 percentile)		
1A Good Quality	<ul style="list-style-type: none"> (i) Dissolved oxygen saturation greater than 80% (ii) Biochemical oxygen demand not greater than 3 mg/l (iii) Ammonia not greater than 0.4 mg/l (iv) Where the water is abstracted for drinking water, it complies with requirements for A2* water (v) Non-toxic to fish in EIFAC terms (or best estimates if EIFAC figures not available) 	<ul style="list-style-type: none"> (i) Average BOD probably not greater than 1.5 mg/l (ii) Visible evidence of pollution should be absent 	<ul style="list-style-type: none"> (i) Water of high quality suitable for potable supply abstractions and for all abstractions (ii) Game or other high class fisheries (iii) High amenity value
1B Good Quality	<ul style="list-style-type: none"> (i) DO greater than 60% saturation (ii) BOD not greater than 5 mg/l (iii) Ammonia not greater than 0.9 mg/l (iv) Where water is abstracted for drinking water, it complies with the requirements for A2* water (v) Non-toxic to fish in EIFAC terms (or best estimates if EIFAC figures not available) 	<ul style="list-style-type: none"> (i) Average BOD probably not greater than 2 mg/l (ii) Average ammonia probably not greater than 0.5 mg/l (iii) Visible evidence of pollution should be absent (iv) Waters of high quality which cannot be placed in Class 1A because of the high proportion of high quality effluent present or because of the effect of physical factors such as canalisation, low gradient or eutrophication (v) Class 1A and Class 1B together are essentially the Class 1 of the River Pollution Survey (RPS) 	<ul style="list-style-type: none"> Water of less high quality than Class 1A but usable for substantially the same purposes
2 Fair Quality	<ul style="list-style-type: none"> (i) DO greater than 40% saturation (ii) BOD not greater than 9 mg/l (iii) Where water is abstracted for drinking water it complies with the requirements for A3* water (iv) Non-toxic to fish in EIFAC terms (or best estimates if EIFAC figures not available) 	<ul style="list-style-type: none"> (i) Average BOD probably not greater than 5 mg/l (ii) Similar to Class 2 of RPS (iii) Water not showing physical signs of pollution other than humic colouration and a little foaming below weirs 	<ul style="list-style-type: none"> (i) Waters suitable for potable supply after advanced treatment (ii) Supporting reasonably good coarse fisheries (iii) Moderate amenity value

3 Poor Quality	(i) DO greater than 10% saturation (ii) Not likely to be anaerobic (iii) BOD not greater than 17 mg/l. This may not apply if there is a high degree of re-aeration	Similar to Class 3 of RPS	Waters which are polluted to an extent that fish are absent, only sporadically present. May be used for low grade industrial abstraction purposes. Considerable potential for further use if cleaned up
----------------	---	---------------------------	---

4 Bad Quality	Waters which are inferior to Class 3 in terms of dissolved oxygen and likely to be anaerobic at times DO greater than 10% saturation	Similar to Class 4 of RPS	Waters which are grossly polluted and are likely to cause nuisance Insignificant watercourses and ditches not usable, where the objective is simply to prevent nuisance developing
---------------	---	---------------------------	---

- Notes
- (a) Under extreme weather conditions (eg flood, drought, freeze-up), or when dominated by plant growth, or by aquatic plant decay, rivers usually in Class 1, 2, and 3 may have BODs and dissolved oxygen levels, or ammonia content outside the stated levels for those Classes. When this occurs the cause should be stated along with analytical results.
 - (b) The BOD determinations refer to 5 day carbonaceous BOD (ATU). Ammonia figures are expressed as NH₄. **
 - (c) In most instances the chemical classification given above will be suitable. However, the basis of the classification is restricted to a finite number of chemical determinands and there may be a few cases where the presence of a chemical substance other than those used in the classification markedly reduces the quality of the water. In such cases, the quality classification of the water should be down-graded on the basis of biota actually present, and the reasons stated.
 - (d) EIFAC (European Inland Fisheries Advisory Commission) limits should be expressed as 95 percentile limits.

EEC category A2 and A3 requirements are those specified in the EEC Council directive of 16 June 1975 concerning the Quality of Surface Water intended for Abstraction of Drinking Water in the Member State.

** Ammonia Conversion Factors

(mg NH₄/l to mg N/l)

Class 1A	0.4 mg NH ₄ /l = 0.31 mg N/l
Class 1B	0.9 mg NH ₄ /l = 0.70 mg N/l
	0.5 mg NH ₄ /l = 0.39 mg N/l

NWC RIVER CLASSIFICATION SYSTEM

CRITERIA USED BY NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION FOR NON-METALLIC DETERMINANDS

River Class	Quality Criteria
1A	Dissolved oxygen % saturation greater than 80% BOD (ATU) not greater than 3 mg/l O Total ammonia not greater than 0.31 mg/l N Non-ionised ammonia not greater than 0.021 mg/l N Temperature not greater than 21.5 C pH greater than 5.0 and less than 9.0 Suspended solids not greater than 25 mg/l
1B	Dissolved oxygen % saturation greater than 60% BOD (ATU) not greater than 5 mg/l O Total ammonia not greater than 0.70 mg/l N Non-ionised ammonia not greater than 0.021 mg/l N Temperature not greater than 21.5 C pH greater than 5.0 and less than 9.0 Suspended solids not greater than 25 mg/l
2	Dissolved oxygen & saturation greater than 40% BOD (ATU) not greater than 9 mg/l O Total ammonia not greater than 1.56 mg/l N Non-ionised ammonia not greater than 0.021 mg/l N Temperature not greater than 28 C pH greater than 5.0 and less than 9.0 Suspended solids not greater than 25 mg/l
3	Dissolved oxygen % saturation greater than 10% BOD (ATU) not greater than 17 mg/l O
4	Dissolved oxygen % saturation not greater than 10% BOD (ATU) greater than 17 mg/l O

STATISTICS USED BY NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION

Determinand	Statistic
Dissolved oxygen	5 percentile
BOD (ATU)	95 percentile
Total ammonia	95 percentile
Non-ionised ammonia	95 percentile
Temperature	95 percentile
pH	5 percentile
Suspended solids	95 percentile
	arithmetic mean

NWC RIVER CLASSIFICATION SYSTEM

CRITERIA USED BY NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION FOR METALLIC DETERMINANDS

SOLUBLE COPPER

Total Hardness (mean) mg/l CaCO ₃	Statistic	Soluble Copper*	
		Class 1 ug/l Cu	Class 2
0 - 10	95 percentile	< = 5	> 5
10 - 50	95 percentile	< = 22	> 22
50 - 100	95 percentile	< = 40	> 40
100 - 300	95 percentile	< = 112	> 112

* Total copper is used for classification until sufficient data on soluble copper can be obtained.


TOTAL ZINC

Total Hardness (mean) mg/l CaCO ₃	Statistic	Total Zinc		
		Class 1 ug/l Zn	Class 2	Class 3
0 - 10	95 percentile	< = 30	< = 300	> 300
10 - 50	95 percentile	< = 200	< = 700	> 700
50 - 100	95 percentile	< = 300	< = 1000	> 1000
100 - 300	95 percentile	< = 500	< = 2000	> 2000

NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION
 1990 RIVER WATER QUALITY CLASSIFICATION
 CATCHMENT: LOOE (16)

1990 Map Position Number	River	Reach upstream of	User Reference Number	National Grid Reference	Reach Length (km)	Distance from source (km)	River Quality Objective	85 RWC Class	86 RWC Class	87 RWC Class	88 RWC Class	89 RWC Class	90 RWC Class
1	EAST LOOE RIVER	VENTON VEOR BRIDGE	R14B005	SX 2304 6577	2.9	2.9	1B	2	2	1B	1B	2	2
2	EAST LOOE RIVER	LOOE MILLS	R14B001	SX 2323 6456	1.0	3.9	1B	2	2	1B	2	2	3
3	EAST LOOE RIVER	LAMELLION MILL	R14B002	SX 2388 6359	1.5	5.4	1B	2	1B	2	2	2	2
4	EAST LOOE RIVER	BELOW LISKEARD STW	R14B008	SX 2422 6280	0.9	6.3	1B	2	3	2	2	2	
5	EAST LOOE RIVER	TRUSSEL BRIDGE	R14B003	SX 2455 6200	0.9	7.2	1B	2	3	2	2	2	2
6	EAST LOOE RIVER	LANDLOOE BRIDGE	R14B006	SX 2500 5950	3.0	10.2	1B	2	3	1B	2	2	2
7	EAST LOOE RIVER	RAILWAY HALT SANDPLACE	R14B004	SX 2483 5715	2.6	12.8	1B	2	3	2	1B	1B	1B
8	DOBWALLS STREAM	TUELMENNA BRIDGE	R14B007	SX 225 651	1.5	1.5	1B						3
	DOBWALLS STREAM	EAST LOOE CONFLUENCE (INFERRED STRETCH)			0.7	2.2	1B						3
9	WEST LOOE RIVER	BOSENT BRIDGE	R14C010	SX 2128 6346	2.0	2.0	1B	1B	1B	3	3	3	3
10	WEST LOOE RIVER	SCAWN MILL BRIDGE	R14C001	SX 2158 6213	1.5	3.5	1B	1B	1B	3	3	3	2
11	WEST LOOE RIVER	CHURCHBRIDGE	R14C002	SX 2193 5858	4.3	7.8	1B	1B	1B	1B	1B	1B	2
12	WEST LOOE RIVER	SOWDEN'S BRIDGE	R14C003	SX 2302 5556	3.7	11.5	1B	1B	3	2	1B	2	2
	WEST LOOE RIVER	NORMAL TIDAL LIMIT (INFERRED STRETCH)			0.6	12.1	1B	1B	3	2	1B	2	2
13	COLDRINNICK STREAM	TREGARRICK MILL BRIDGE	R14C011	SX 2058 5713	3.2	3.2	1B	1B		2	1B	2	1B
	COLDRINNICK STREAM	WEST LOOE CONFLUENCE (INFERRED STRETCH)			1.8	5.0	1B	1B		2	1B	2	1B
14	CONNON STREAM	ABOVE WASTE DISPOSAL SITE	R14C005	SX 1880 6259	1.3	1.3	1B	1B	2	4	4	4	3
15	CONNON STREAM	TREVILLIS WOOD	R14C006	SX 1962 6178	1.4	2.7	1B	1B	2	2	2	2	3
16	CONNON STREAM	HERODSFOOT BRIDGE	R14C008	SX 2140 6042	2.5	5.2	1B	1B	2	2	1B	1B	1B
	CONNON STREAM	WEST LOOE CONFLUENCE (INFERRED STRETCH)			0.1	5.3	1B	1B	2	2	1B	1B	1B
17	POLPERRO RIVER	POLPERRO	R14A001	SX 2088 5097	6.8	6.8	1B	1B	1B			2	2
	POLPERRO RIVER	NORMAL TIDAL LIMIT (INFERRED STRETCH)			0.2	7.0	1B	1B	1B			2	2


Looe Catchment Water Quality - 1990



NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION
 1990 RIVER WATER QUALITY CLASSIFICATION
 CALCULATED DETERMINED STATISTICS USED FOR QUALITY ASSESSMENT
 CRUICMENT: LOOE (16)

River	Reach upstream of	User Ref. Number	90 BAC Class	Calculated Determined Statistics used for Quality Assessment																			
				pH Lower Class 5tile		pH Upper Class 95tile		Temperature Class 95tile		DO (%) Class 5tile		BOD (MU) Class 95tile		Total Ammonia Class 95tile		Union. Ammonia Class 95tile		S.Solids Class Mean		Total Copper Class 95tile		Total Zinc Class 95tile	
EAST LOOE RIVER	VENISEN VBOUR BRIDGE	[R14B005]	2	1A	7.0	1A	7.8	1A	16.9	1A	82.2	1B	4.6	1B	0.396	1A	0.010	1A	21.8	2	59.0	1A	294.5
EAST LOOE RIVER	LOOE MILLS	[R14B001]	3	1A	7.1	1A	7.7	1A	16.0	1A	80.2	2	7.0	1B	0.454	1A	0.010	3	26.7	1A	16.0	1A	44.0
EAST LOOE RIVER	LAMELLETON MILL	[R14B002]	2	1A	7.1	1A	7.9	1A	17.8	1B	70.2	2	5.2	1B	0.435	1A	0.010	1A	21.0	1A	26.4	1A	44.0
EAST LOOE RIVER	TRUSSELL BRIDGE	[R14B003]	2	1A	7.2	1A	7.6	1A	16.4	1B	70.0	2	7.2	1B	0.635	1A	0.010	1A	24.5	1A	19.4	1A	68.0
EAST LOOE RIVER	LANDLOOE BRIDGE	[R14B006]	2	1A	7.3	1A	7.8	1A	16.4	1A	82.6	2	5.1	1B	0.320	1A	0.010	1A	23.6	1A	16.7	1A	38.5
EAST LOOE RIVER	BRIDWAY HUNT SPURPLACE	[R14B004]	1B	1A	7.3	1A	7.9	1A	16.6	1A	81.2	1B	3.4	1A	0.223	1A	0.010	1A	15.5	1A	8.7	1A	48.0
DOEWALLS STREAM	TUBEMENNA BRIDGE	[R14B007]	3	1A	6.6	1A	7.7	1A	17.5	1A	87.0	2	5.5	1B	0.480	1A	0.010	3	30.8	1A	13.0	1A	47.0
WEST LOOE RIVER	ROBERT BRIDGE	[R14C010]	3	1A	7.0	1A	7.8	1A	15.0	1B	67.0	3	9.3	3	2.598	1A	0.010	3	46.7	1A	14.0	1A	54.0
WEST LOOE RIVER	SCOWS MILL BRIDGE	[R14C001]	2	1A	7.1	1A	8.3	1A	16.2	1A	81.4	2	6.5	2	1.045	1A	0.010	1A	11.8	1A	11.0	1A	29.0
WEST LOOE RIVER	CHURCHERIDGE	[R14C002]	2	1A	7.1	1A	7.9	1A	15.8	1A	83.3	2	5.2	2	1.286	1A	0.010	1A	24.3	1A	16.0	1A	42.8
WEST LOOE RIVER	SOMDEN'S BRIDGE	[R14C003]	2	1A	7.1	1A	7.8	1A	15.4	1B	78.0	1B	4.0	1A	0.282	1A	0.010	1A	18.0	2	297.8	2	722.4
COLDKIRKICK STREAM	INGLAROCK MILL BRIDGE	[R14C011]	1B	1A	6.8	1A	7.8	1A	15.8	1B	78.3	1B	4.2	1A	0.203	1A	0.010	1A	15.0	1A	14.0	1A	53.0
CORNON STREAM	ABOVE WASTE DISPOSAL SITE	[R14C005]	3	1A	6.5	1A	8.2	1A	14.8	1B	75.7	2	8.7	3	3.050	1A	0.019	3	94.6	2	320.4	2	759.0
CORNON STREAM	TREVILLIS WOOD	[R14C006]	3	1A	6.5	1A	7.7	1A	15.4	1B	71.0	1B	4.7	3	1.670	1A	0.010	1A	24.5	1A	11.7	1A	45.5
CORNON STREAM	HERODSFOOT BRIDGE	[R14C008]	1B	1A	6.8	1A	8.1	1A	15.2	1B	71.9	1B	3.2	1B	0.595	1A	0.010	1A	17.1	1A	9.9	1A	30.6
FOLFERRO RIVER	FOLFERRO	[R14A001]	2	1A	7.4	1A	8.2	1A	15.9	1A	81.4	2	7.8	1A	0.132	1A	0.010	1A	18.5	1A	8.0	1A	31.9

Looe Catchment Water Quality - 1990



NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION

1990 RIVER WATER QUALITY CLASSIFICATION

NUMBER OF SAMPLES (N) AND NUMBER OF SAMPLES EXCEEDING QUALITY STANDARD (P)

CATCHMENT: LOCE (16)

River	Reach upstream of	User Ref. Number	pH Lower		pH Upper		Temperature		DO (%)		BOD (RTU)		Total Ammonia		Union. Ammonia		S.Solids		Total Copper		Total Zinc	
			N	P	N	P	N	P	N	P	N	P	N	P	N	P	N	P	N	P	N	P
EAST LOCE RIVER	VENION VEOR BRIDGE	R14B005	25	-	25	-	25	-	25	-	25	1	25	-	21	-	25	2	24	1	24	1
EAST LOCE RIVER	LOCE MILLS	R14B001	32	-	32	-	32	-	32	-	32	1	32	-	31	-	32	4	24	-	24	-
EAST LOCE RIVER	LAMELLION MILL	R14B002	24	-	24	-	24	-	23	-	24	1	24	-	24	-	24	4	23	-	23	-
EAST LOCE RIVER	TRUSSEL BRIDGE	R14B003	32	-	32	-	29	-	29	-	32	2	32	1	28	-	32	3	28	-	28	-
EAST LOCE RIVER	LANDLOCE BRIDGE	R14B006	26	-	26	-	24	-	23	-	26	1	26	-	24	-	26	4	26	-	26	-
EAST LOCE RIVER	RAILWAY HALT SANDPLACE	R14B004	32	-	32	-	31	-	31	-	31	-	32	-	31	-	32	4	25	-	25	-
DOBWALLS STREAM	TUELMENNA BRIDGE	R14B007	15	-	15	-	15	-	15	-	15	1	15	-	14	-	15	3	14	-	14	-
WEST LOCE RIVER	BOSENT BRIDGE	R14C010	24	-	24	-	23	-	23	-	24	1	24	5	23	-	24	4	15	-	15	-
WEST LOCE RIVER	SCAWN MILL BRIDGE	R14C001	24	-	24	-	23	-	23	-	24	1	24	2	21	-	24	3	23	-	23	-
WEST LOCE RIVER	CHURCHBRIDGE	R14C002	25	-	25	-	24	-	24	-	25	1	25	1	21	-	25	5	24	-	24	-
WEST LOCE RIVER	SCODEN'S BRIDGE	R14C003	34	-	34	-	32	-	33	-	34	1	34	-	29	-	34	4	27	1	27	1
COLDRINNICK STREAM	TRIGARRICK MILL BRIDGE	R14CD11	24	-	24	-	24	-	24	-	24	-	24	-	23	-	24	4	14	-	14	-
CONNON STREAM	ABOVE WASTE DISPOSAL SITE	R14CD05	25	-	25	-	26	-	26	-	25	3	25	8	21	-	25	4	26	1	26	1
CONNON STREAM	TRIVILLIS WOOD	R14CD06	25	-	25	-	23	-	23	-	25	1	25	4	20	-	25	3	25	-	25	-
CONNON STREAM	HENDSPOOT BRIDGE	R14CD08	32	-	32	-	32	-	32	-	32	-	32	1	31	-	32	4	28	-	28	-
FOLFERRO RIVER	FOLFERRO	R14A001	24	-	24	-	24	-	23	-	24	2	24	-	23	-	24	3	21	-	21	-

NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION

1990 RIVER WATER QUALITY CLASSIFICATION

PERCENTAGE EXCEEDENCE OF DETERMINAND STATISTICS FROM QUALITY STANDARDS

CATCHMENT: LOOE (16)

River	Reach upstream of	User Ref. Number	PERCENTAGE EXCEEDENCE OF STATISTIC FROM QUALITY STANDARD									
			pH Lower	pH Upper	Temperature	DO (%)	BOD (ATU)	Total Ammonia	Un-ionised Ammonia	Suspended Solids	Total Copper	Total Zinc
EAST LOOE RIVER	VENTON VEOR BRIDGE	R14B005	-	-	-	-	-	-	-	-	48	-
EAST LOOE RIVER	LOOE MILLS	R14B001	-	-	-	-	41	-	-	7	-	-
EAST LOOE RIVER	LAMELLION MILL	R14B002	-	-	-	-	4	-	-	-	-	-
EAST LOOE RIVER	TRUSSEL BRIDGE	R14B003	-	-	-	-	45	-	-	-	-	-
EAST LOOE RIVER	LANDLOOE BRIDGE	R14B006	-	-	-	-	2	-	-	-	-	-
EAST LOOE RIVER	RAILWAY HALT SANDPLACE	R14B004	-	-	-	-	-	-	-	-	-	-
DOBWALLS STREAM	TUELMERNA BRIDGE	R14B007	-	-	-	-	10	-	-	23	-	-
WEST LOOE RIVER	BOSENT BRIDGE	R14C010	-	-	-	-	85	271	-	87	-	-
WEST LOOE RIVER	SCAWN MILL BRIDGE	R14C001	-	-	-	-	29	49	-	-	-	-
WEST LOOE RIVER	CHURCHBRIDGE	R14C002	-	-	-	-	4	84	-	-	-	-
WEST LOOE RIVER	SOWDEN'S BRIDGE	R14C003	-	-	-	-	-	-	-	-	645	141
COLDRINNICK STREAM	TREGARRICK MILL BRIDGE	R14C011	-	-	-	-	-	-	-	-	-	-
CONNON STREAM	ABOVE WASTE DISPOSAL SITE	R14C005	-	-	-	-	74	336	-	278	701	153
CONNON STREAM	TREVILLIS WOOD	R14C006	-	-	-	-	-	139	-	-	-	-
CONNON STREAM	HERODSFOOT BRIDGE	R14C008	-	-	-	-	-	-	-	-	-	-
POLPERRO RIVER	POLPERRO	R14A001	-	-	-	-	55	-	-	-	-	-

NATIONAL RIVERS AUTHORITY - SOUTH WEST REGION
 IDENTIFICATION OF POSSIBLE CAUSES OF NON-COMPLIANCE WITH RQO
 CATCHMENT: LOOE (16)

* = WORK ALREADY IN HAND

1990 Map Position Number	River	Reach upstream of	User Reference Number	Reach Length (km)	Possible causes of non-compliance
1	EAST LOOE RIVER	VENTON VEOR BRIDGE	R14B005	2.9	LAND RUN-OFF
2	EAST LOOE RIVER	LOOE MILLS	R14B001	1.0	LAND RUN-OFF, INDUSTRIAL DISCHARGE
3	EAST LOOE RIVER	LAMELLION MILL	R14B002	1.5	LAND RUN-OFF, FARMING ACTIVITIES
5	EAST LOOE RIVER	* TRUSSEL BRIDGE	R14B003	0.9	LAND RUN-OFF, SEWAGE TREATMENT WORKS
6	EAST LOOE RIVER	LANDLOOE BRIDGE	R14B006	3.0	LAND RUN-OFF
8	DOBMAIS STREAM	TUELMENNA BRIDGE	R14B007	1.5	LAND RUN-OFF, CHINA CLAY DISCHARGE
9	WEST LOOE RIVER	BOSENT BRIDGE	R14C010	2.0	LAND RUN-OFF, SEWAGE TREATMENT WORKS
10	WEST LOOE RIVER	SCAWN MILL BRIDGE	R14C001	1.5	LAND RUN-OFF, SEWAGE TREATMENT WORKS
11	WEST LOOE RIVER	CHURCHBRIDGE	R14C002	4.3	LAND RUN-OFF, SEPTIC TANKS
12	WEST LOOE RIVER	SOWDEN'S BRIDGE	R14C003	3.7	MINING
14	CORNON STREAM	ABOVE WASTE DISPOSAL SITE	R14C005	1.3	LAND RUN-OFF, FARMING ACTIVITIES, WASTE DISPOSAL SITE
15	CORNON STREAM	* TREVILLIS WOOD	R14C006	1.4	WASTE DISPOSAL SITE, LAND RUN-OFF
17	POLPERRO RIVER	POLPERRO	R14A001	6.8	LAND RUN-OFF, URBANISATION, CARALISATION